

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 December 2007 (27.12.2007)

PCT

(10) International Publication Number
WO 2007/149832 A2

(51) International Patent Classification:
A61M 31/00 (2006.01)

(74) Agents: **JOHNSON, Philip S.** et al.; One Johnson & Johnson Plaza, New Brunswick, New Jersey 08933 (US).

(21) International Application Number:
PCT/US2007/071520

(22) International Filing Date: 19 June 2007 (19.06.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/805,378 21 June 2006 (21.06.2006) US
11/759,327 7 June 2007 (07.06.2007) US

(71) Applicant (for all designated States except US): **JOHNSON & JOHNSON VISION CARE, INC.** [US/US]; 7500 Centurian Parkway, Suite 100, Jacksonville, Florida 32256 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BORGIA, Maureen J.** [US/US]; 111 North Clark Avenue, Somerville, New Jersey 08876 (US). **CHAOUK, Hassan** [AU/US]; 7944 Hampton Park Boulevard East, Jacksonville, Florida 32256 (US). **CUI, Han** [CN/US]; 7 Liberty Ridge Road, Basking Ridge, New Jersey 07920 (US). **LI, Zhigang** [US/US]; 83 Fisher Drive, Hillsborough, New Jersey 08844 (US). **NATHAN, Aruna** [IN/US]; 16 Woodward Drive, Bridgewater, New Jersey 08807 (US). **TREZZA II, Michael J** [US/US]; 4 Pheasant Run Road, Great Meadows, New Jersey 07838 (US). **LAREDO, Walter** [US/US]; 3901 Es-tancia Way, Fort Worth, Texas 76108 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PUNCTAL PLUGS FOR THE DELIVERY OF ACTIVE AGENTS



(57) Abstract: The invention provides punctal plugs for the delivery of active agent to one or both of the tear fluid of the eye and to the nasolacrimal duct. The plugs of the invention have a body, a reservoir contained within the body, and optionally a collarette. The reservoir has at least one opening and contains a polymeric material and at least one active agent.

WO 2007/149832 A2

PUNCTAL PLUGS FOR THE DELIVERY OF ACTIVE AGENTS

Field of the Invention

5 The present invention relates to devices suitable for delivering substances to one or more of the eye, nose and throat. In particular, the invention relates to punctal plugs for delivery of at least one active agent.

Related Applications

10 This application claims priority from provisional application United States Serial No. 60/805,378 filed on June 21, 2006.

Background of the Invention

 Human tears are secreted by the lacrimal gland and flow across the surface of
15 the eye to a shallow pool, known as the lacrimal lake, located where the eyelids come together at their inner ends. From there, the tears drain through small openings in each of the upper and lower eyelids, termed the superior lacrimal punctum and the inferior lacrimal punctum, respectively. From the superior and inferior puncta, the tears pass into each of the superior and inferior lacrimal
20 canaliculus, respectively, which are duct-like pathways that lead to the lacrimal sac. The lacrimal sac is the superior, expanded portion of the nasolacrimal duct, which drains tears into the nasal system. Active agents can thus be delivered to the nose and throat through the lacrimal canaliculi, which lead into the nasolacrimal duct.

25 Active agents frequently are administered to the eye for the treatment of ocular diseases and disorders. Conventional means for delivering active agents to the eye involve topical application to the surface of the eye. The eye is uniquely suited to topical administration because, when properly constituted, topically applied active agents can penetrate through the cornea and rise to therapeutic concentration
30 levels inside the eye. Active agents for ocular diseases and disorders may be administered orally or by injection, but such administration routes are

disadvantageous in that, in oral administration, the active agent may reach the eye in too low a concentration to have the desired pharmacological effect and their use is complicated by significant, systemic side effects, while injections pose the risk of infection.

The majority of ocular active agents are currently delivered topically using eye drops which, though effective for some applications, are inefficient. When a drop of liquid is added to the eye, it overfills the conjunctival sac, the pocket between the eye and the lids, causing a substantial portion of the drop to be lost due to overflow of the lid margin onto the cheek. In addition, a substantial portion of the drop that remains on the ocular surface is drained into the lacrimal puncta, diluting the concentration of the drug.

Brief Description of the Drawings

Figure 1 is a sectional view of a punctal plug having a body 10 with an enlarged segment 12 and a reservoir 15 within the body 10 that contains a polymeric material 11 that contains active agent 18. The reservoir 15 has an opening 13 through which the active agent 18 is released.

Figure 2 is a sectional view of a punctal plug having a body 20 with an enlarged segment 22 and a reservoir 25 within the body 20 that contains a polymeric material 21 that contains active agent 28. The reservoir 25 has an opening 23 through which the active agent 28 is released.

Figure 3 is a sectional view of a punctal plug having a body 30 with an enlarged segment 32 and a reservoir 35 within the body 30 that contains a polymeric material 31 that contains active agent 38. The reservoir 35 has two openings 33 and 33' through which the active agent 38 is released.

Figure 4 is a sectional view of a punctal plug having a body 40 with an enlarged segment 42, a reservoir 45 within the body 40 that contains a polymeric material 41 that contains active agent 48, and a collarette 44. The reservoir 45 has an opening 43 through which the active agent 48 is released.

Figure 5 is a sectional view of a punctal plug having a body 50 with an enlarged segment 52, a reservoir 55 within the body 50 that contains a polymeric material 51 that contains active agent 58, and a collarette 54. The reservoir 55 has an opening 53 through which the active agent 58 is released.

Figure 6 is a sectional view of a punctal plug having a body 60 with an enlarged segment 62, a reservoir 65 within the body 60 that contains a polymeric material 61 that contains active agent 68, and a collarette 64. The reservoir 65 has two openings 63 and 63' through which the active agent 68 is released.

Figure 7 is a sectional view of a punctal plug having a body 70 made of a flexible polymeric material, a reservoir 75 within the body 70 that is coterminous with the body 70, and a collarette 74. The reservoir 75 contains a polymeric material 71 that contains active agent 78, and the reservoir 75 has an opening 73 through which the active agent 78 is released.

Figure 8 is a sectional view of a punctal plug having a body 80 made of a flexible polymeric material, a reservoir 85 within the body 80 that is coterminous with the body 80, and a collarette 84. The reservoir 85 contains a polymeric material 81 that contains active agent 88, and the reservoir 85 has two openings 83 and 83' through which the active agent 88 is released.

Figure 9 is a three-dimensional view of the punctal plug depicted two-dimensionally in Figure 4 having a body 90 with an enlarged segment 92, a reservoir 95 within the body 90 that contains a polymeric material 91 that contains active

agent 98, and a collarette 94. The reservoir 95 has an opening 93 through which the active agent 98 is released.

5 Figure 10 is a sectional view of a punctal plug having a body 100 with an enlarged segment 102, a reservoir 105 within the body 100, a collarette 104, and a tapering, rounded end 103. The reservoir 105 contains a polymeric material 101 that contains active agent 108, and the reservoir 105 has openings 103 through which the active agent 108 is released.

10

Figure 11 is a graph depicting the active agent release profile for the punctal plug of Example 2.

15 Figure 12 is a graph depicting the active agent release profile for the punctal plug of Example 3.

Detailed Description of the Invention and Illustrative Embodiments

The present invention provides punctal plugs that can be used to deliver active agents to one or both of the nasolacrimal duct and to the tear fluid of the eye.

20 In one embodiment, the invention provides punctal plugs comprising, consisting essentially of, and consisting of: a body having a first end and a second end; a lateral surface extending between the two ends; a reservoir contained within the body wherein the reservoir comprises, consists essentially of and consists of at least one opening and contains a material that comprises, consists essentially of and consists
25 of at least one active agent; and wherein the body is impermeable to the active agent.

Referring to Figure 1, punctal plug body 10 has a reservoir 15 that contains at least one opening 13 and active agent 18 is released through opening 13, for example, when the active agent-containing material 11, preferably a polymeric material, dissolves, degrades, or the active agent 18 simply diffuses or is released
30 from material 11, depending upon the nature of the material. The opening through

which the active agent is released from the plug may be located at a first end, as for example in Figure 1, a second end as for example in Figure 2, or both the first and second ends of the plug body as for example shown in Figure 3, or along the lateral
5 surface thereof. Preferably, the opening is located at one or both of the first and second ends. In particular embodiments of the invention, for example as shown in Figure 3, the punctal plug contains an enlarged segment 32 of the body 30 that is of a suitable size and shape for securing the punctal plugs in the lacrimal canaliculus.

10 For delivery of an active agent into the tear fluid of the eye, a punctal plug is inserted into a lacrimal canaliculus and the active agent is released into the tear fluid of the eye. Referring to Figure 4, for delivery into the tear fluid, a collarette 44 is preferably provided on body 40 of the punctal plug and, when the punctal plug is inserted into the lacrimal canaliculus, the collarette 44 rests on the exterior of the
15 lacrimal punctum. For delivery of active agent into the nasolacrimal duct, a punctal plug is inserted, preferably deeply, into the lacrimal canaliculus and the active agent is released into the nasolacrimal duct.

As used herein, the term “punctal plug” refers to a device of a size and shape
20 suitable for insertion into the inferior or superior lacrimal canaliculus of the eye through the inferior or superior lacrimal punctum.

As used herein, the term “active agent” refers to an agent capable of treating, inhibiting, or preventing a disorder or a disease. Exemplary active agents include,
25 without limitation, pharmaceuticals and nutraceuticals. Preferred active agents are capable of treating, inhibiting, or preventing a disorder or a disease of one or more of the eye, nose and throat.

As used herein, the phrase “a material that is at least partially water-soluble” refers to a material that exhibits a level of solubility in water sufficient to result in detectable dissolution of the material upon exposure to an aqueous environment.

5

As used herein, the phrase “a material that is biodegradable” refers to a material that degrades to a detectable degree upon exposure to biologically active substances typically present in mammals.

10

As used herein, the phrase “a material that is insoluble in water” refers to a material that does not dissolve to a substantial degree upon exposure to water.

As used herein, the phrase “a material that is non-biodegradable” refers to a material that does not degrade to a substantial degree upon exposure to biologically
15 active substances typically present in mammals.

As used herein, the phrase “body that is impermeable to active agent” refers to a body through which only an insubstantial amount of active agent can pass.

20

As used herein, the term “polymeric material” refers to a material made of one or more types of polymers that is capable of containing at least one active agent and releasing the active agent, for example, when the polymers dissolve or degrade, when the active agent diffuses from the polymers, or when a pro-drug is used in which the active agent is attached to the polymers and then released by being
25 cleaved from the material.

As used herein, the term “opening” refers to an opening in the body of a punctal plug of a size and shape through which the active agent can pass. Preferably, only the active agent can pass through the opening. The opening, for
30 example, may be a hole covered with a membrane, mesh, grid or it may be

uncovered. The membrane, mesh, or grid may be one or more of porous, semi-porous, permeable, semi-permeable, and biodegradable.

5

As used herein, the phrase “flexible material” refers to a material that is not rigid and that substantially conforms to the surface of whatever object the material contacts.

10

As used herein, the phrase “the reservoir and the body are coterminous” indicates that the reservoir is substantially all of the body. A collarette can be attached to the body when the reservoir and body are coterminous, but the collarette would not be considered to be part of the body.

15

As used herein, the phrase “refilled with active agent” refers to adding any detectable amount of active agent to the reservoir of a punctal plug.

20

The present invention encompasses numerous punctal plugs for the delivery of active agents to one or both of the tear fluid of the eye and to the nasolacrimal duct. The punctal plugs preferably are inserted into the inferior lacrimal canaliculus, the superior lacrimal canaliculus, or both the inferior and superior lacrimal canaliculi. If the punctal plugs are being used to deliver active agents to the tear fluid of the eye, the punctal plugs preferably have a collarette at one end of the body. The collarette is a portion of the punctal plug that extends radially outwardly from one end of the body to a degree sufficient, and having a size and a shape, such that at least a portion of the collarette will extend beyond and be exterior to the lacrimal punctum after insertion of the punctal plug into the lacrimal canaliculus. Typically, the collarette will extend about 0.2 to about 1 mm beyond the plug body. The portion of the punctal plug without the collarette is inserted into one of the inferior lacrimal punctum or the superior lacrimal punctum. Referring to Figure 6, enlarged segment 62 and body 60 is inserted into one of the punctum, and collarette 64 rests

30

against the exterior of the lacrimal punctum and keeps the punctal plug from
slipping down into the lacrimal canaliculus, so that contact between the punctal plug
5 and the tear fluid of the eye is maintained.

If the punctal plugs are being used to deliver active agent to the nasolacrimal
duct, the punctal plugs may not have a collarette so that they may be inserted at a
sufficient depth within one or both of the lacrimal canaliculi such that the active
10 agent is released into the lacrimal sac. In Figures 1 through 3 are depicted examples
of punctal plugs useful for delivery of an active agent into the nasolacrimal duct.

The numerous punctal plugs of the invention each have various features and
advantages. For example, certain punctal plugs have a body with a first end, a
15 second end, and a lateral surface extending between the two ends. The lateral
surface preferably has an outer diameter that is substantially circular in shape and,
thus, the body preferably has a cylindrical shape. A portion of the lateral surface of
certain of the punctal plugs preferably has an outer diameter that is greater than the
outer diameter of the remainder of the lateral surface. With reference to Figure 2,
20 the enlarged portion 22 of the lateral surface anchors or secures the punctal plugs in
the lacrimal canaliculus. The enlarged portion can be any size or shape, and can be
present on any part of the lateral surface, so long as the enlarged portion at least
partially anchors the punctal plug in the lacrimal canaliculus. Preferably, the
enlarged portion is at one end of the plug. Conveniently, the enlarged portion may
25 take the shape of an inverted triangle having a flattened apex, as shown in Figure 1,
may have an untapered, body rounded at the end, or may have a tapered shape at one
end with a rounded point as shown in Figure 10. One ordinarily skilled in the art
will recognize that any of a wide variety of shapes are possible.

30 The body of the punctal plugs of the invention may take any shape and size,
Preferably, the body is in the shape of an elongated cylinder. The body will be

about 0.8 to about 5 mm in length, preferably about 1.2 to about 2.5 mm in length.
The width of the body will be about 0.2 to about 3, preferably 0.3 to about 1.5 mm.

5

The body of the plug may be wholly or partially transparent or opaque.
Optionally, the body may include a tint or pigment that makes the plug easier to see
when it is placed in a punctum.

10 The body of the punctal plugs may be made of any suitable biocompatible
material including, without limitation, silicone, silicone blends, silicone co-
polymers, such as, for example, hydrophilic monomers of
polyhydroxyethylmethacrylate ("pHEMA"), polyethylene glycol,
polyvinylpyrrolidone, and glycerol, and silicone hydrogel polymers such as, for
15 example, those described in U.S. Patent Nos. 5,962,548, 6,020,445, 6,099,852,
6,367,929, and 6,822,016, incorporated herein in their entireties by reference. Other
suitable biocompatible materials include, for example: polyurethane;
polymethylmethacrylate; poly(ethylene glycol); poly(ethylene oxide);
poly(propylene glycol); poly(vinyl alcohol); poly(hydroxyethyl methacrylate);
20 poly(vinylpyrrolidone) ("PVP"); polyacrylic acid; poly(ethylloxazoline);
poly(dimethyl acrylamide); phospholipids, such as, for example, phosphoryl choline
derivatives; polysulfobetains; acrylic esters, polysaccharides and carbohydrates,
such as, for example, hyaluronic acid, dextran, hydroxyethyl cellulose, hydroxyl
propyl cellulose, gellan gum, guar gum, heparan sulfate, chondritin sulfate, heparin,
25 and alginate; proteins such as, for example, gelatin, collagen, albumin, and
ovalbumin; polyamino acids; fluorinated polymers, such as, for example,
polytetrafluoroethylene ("PTFE"), polyvinylidene fluoride ("PVDF"),
and teflon; polypropylene; polyethylene; nylon; and ethylene vinyl alcohol
("EVA").

30

The surface of the plug body may be wholly or partially coated. The coating may provide one or more of lubriciousness to aid insertion, muco-adhesiveness to improve tissue compatibility, and texture to aid in anchoring the plug within the punctum. Examples of suitable coatings include, without limitation, gelatin, collagen, hydroxyethyl methacrylate, PVP, PEG, heparin, chondroitin sulphate, hyaluronic acid, synthetic and natural proteins, and polysaccharides, thiomers, thiolated derivatives of polyacrylic acid and chitosan, polyacrylic acid, carboxymethyl cellulose and the like and combinations thereof.

10

Certain embodiments of the punctal plugs of the invention have a body made of a flexible material that conforms to the shape of whatever it contacts. Optionally, the plug may have a collarette formed of either a less flexible material than that of the body or material that too conforms to the shape of whatever it contacts. When a punctal plug having both a flexible body and a less flexible collarette is inserted into the lacrimal canaliculus, the collarette rests on the exterior of the lacrimal punctum and the body of the punctal plug conforms to the shape of the lacrimal canaliculus. The reservoir and the body of such punctal plugs are preferably coterminous. That is, the reservoir of such punctal plugs preferably make up the entirety of the body, except for the collarette.

20

In embodiments in which one or both of a flexible body and collarette are used, the flexible body and flexible collarette can be made of materials that include, without limitation, nylon, polyethylene terephthalate ("PET"), polybutylene terephthalate ("PBT"), polyethylenec, polyurethane, silicone, PTFE, PVDF, and polyolefins. Punctal plugs made of nylon, PET, PBT, polyethylene, PVDF, or polyolefins are typically manufactured for example and without limitation, extrusion, injection molding, or thermoforming. Punctal plugs made of latex, polyurethane, silicone, or PTFE are typically manufactured using solution casting processes.

30

The punctal plugs of the invention contain a reservoir within the body, and the reservoir contains an active agent-containing material. The material may be any material that is compatible with the active agent or agents to be delivered by the plug and is capable of releasing the active agent in the desired manner, for example by dissolving or degrading of the material or diffusion of the active agent from the material. Any number of material may be used as the active agent-containing material including, without limitation, polymeric materials, both naturally occurring and synthetic, non-polymeric materials including, without limitation, glasses and clays, organic materials, inorganic materials including, without limitation, porous ceramics, lipids, waxes and the like and combinations thereof. Preferably, the active agent containing-material is a polymeric material, in which at least one active agent is disposed on, dispersed throughout, or otherwise contained. The body is preferably impermeable to the active agent, and the reservoir has at least one opening through which the active agent is released.

The body has one or more openings communicating with the reservoir at a first end, as shown in Figure 4, a second end as shown in Figure 5, both the first and second ends of the body as shown in Figure 6, or at another location on the body. In particular embodiments of the invention, when such punctal plugs are inserted into the lacrimal canaliculus and have opening at the end of the body facing the eye, the active agent is released into the tear fluid of the eye. Alternatively, if the plug has an opening in the end of the body facing the nasolacrimal duct, the active agent is released into the nasolacrimal duct. In those embodiments in which the plug has opening at the end of the body facing the eye and another opening at the end of the body facing the nasolacrimal duct, the active agent is released into both the tear fluid of the eye and the nasolacrimal duct. For those punctal plugs with a collarette, the opening of such punctal plugs is preferably located within the collarette, preferably the central portion of the collarette. When such punctal plugs are inserted into the

lacrimal canaliculus, the opening faces the eye, and the active agent is released into the tear fluid of the eye.

- 5 The size of the opening will be from about 1 nm to about 2.5 mm and preferably about 0.15 mm to about 0.8 mm. Instead of one large opening at any one location, multiple small openings may be used.

Processes for manufacturing the punctal plugs useful in the invention are
10 well known. Typically, the plugs are manufactured by injection molding, cast molding, transfer molding or the like. Preferably, the reservoir is filled with one or both of at least one active agent and the active agent-containing material subsequent to the manufacture of the plug. Additionally, one or more excipients may be combined with the active agent alone or in combination with the polymeric material.

15

Depending upon the active agent-containing material selected, the active agent can be released from the material almost immediately, or the active agent can be released in a sustained manner over a desired period of time. For example, a polymeric material may be used that is composed of one or more polymers that are
20 at least partially soluble in water. When such a polymeric material is exposed to the aqueous environment of the lacrimal canaliculus or the tear fluid, it preferably will dissolve and release the active agent as it dissolves. The solubility in water of the one or more polymers from which the polymeric material is made typically will be directly proportional to its rate of dissolution. Suitable polymers that are at least
25 partially soluble in water include, without limitation, poly(ethylene glycol); poly(ethylene oxide); poly(propylene glycol); poly(vinyl alcohol); poly(hydroxyethyl methacrylate); poly(vinylpyrrolidone); polyacrylic acid; poly(ethyloxazoline); poly(dimethyl acrylamide); phospholipids, such as, for example, phosphoryl choline derivatives; polysulfobetains; polysaccharides and
30 carbohydrates, including, without limitation, hyaluronic acid, dextran, hydroxyethyl

cellulose, hydroxyl propyl cellulose, gellan gum, guar gum, heparan sulfate, chondritin sulfate, heparin, and alginate; proteins such as, for example, gelatin, collagen, albumin, and ovalbumin; and polyamino acids. The polymeric materials in this list can typically be copolymerized or blended with one or both of hydrophobic polymers and monomers.

As an alternative, a non-polymeric material including, without limitation, a lipid, wax, or inorganic material may be used. Suitable non-polymeric materials include, without limitation, lanolin, paraffin, sorbates, lecithin, vitamin A, D, and E, glycerine, sorbitol, mannitol, stearates, fatty acids, lutein, zeaxanthin, taurine, glutathione and the like.

Alternatively, the active agent-containing material can be one or more biodegradable polymers that partially or wholly chemically degrade upon exposure to, for example, biologically active substances typically present in mammals. The biodegradable materials are preferably hydrolyzable under *in vivo* conditions. Biodegradation may occur more slowly than dissolution, and the material can thus compose one or more biodegradable polymers if slower, more sustained release of the active agent is desired.

Suitable biodegradable polymers include, without limitation, polymers and oligomers of glycolide, lactide, lactones, and other hydroxy acids, and other biologically degradable polymers that yield materials that are non-toxic or present as normal metabolites in the body. Preferred poly(alpha-hydroxy acids) are poly(glycolic acid), poly(2-dioxanone); poly(DL-lactic acid) and poly(L-lactic acid). Other useful polymers include poly(amino acids), polycarbonates, poly(anhydrides), poly(orthoesters), poly(phosphazines) and poly(phosphoesters). Polylactones including, without limitation, poly(epsilon-caprolactone), poly(delta-caprolactone),

poly(delta- valerolactone) and poly(gamma-butyrolactone are also useful, as are chitosan, alginates, collagen, and gelatin. In particular aspects of the invention, the
5 polymeric material the contains the active agent can comprise a mixture of one or more dissolvable and bio-degradable polymers.

In a preferred embodiment, the active agent –containing material is a polymeric material that is combined with at least one active agent to form one or
10 more fiber or fiber-like structures, the dimensions of which may be substantially the dimensions of the reservoir or smaller than such dimensions, and one or more of the fibers or fiber-like structures are inserted into the reservoir through the opening in the plug body. The fibers or fiber-like structures may be of a size and a shape suitable for insertion into the opening and may be about 0.5 to about 5 mm in length
15 and 0.05 to about 2 mm in diameter. If only one fiber or fiber-like structure is used, preferably, the dimensions of the fiber are such that the fiber fits securely into the reservoir and remains in the reservoir when the plug is in use in a wearer's punctum. Thus, the fiber can be symmetrical or asymmetrical, depending upon the shape of the reservoir. The internal walls of the reservoir may be substantially smooth or
20 may include features that aid in maintaining the fiber within the reservoir including, without limitation, surfaces with grooves, indentations, roughness or the like in the interior walls.

Alternatively, the fiber containing the active agent or agents may be formed
25 and the plug cast around the fiber. As yet another alternative, the fiber and active agent may be dosed into the plug reservoir as a melt and allowed to solidify. As still another alternative, the polymer and active agent may be introduced as a solution. The solution may contain monomers, pre-polymers and the like suitable for cross-linking via one or more of irradiation, redox, and thermal radical polymerization.
30 As yet another alternative, the fiber may simply be soaked in the active agent before or after insertion in the plug.

Preferably the fiber or fiber-like structures are composed of a polymeric material and more preferably a polymeric material that is polycaprolactone, still more preferably poly(epsilon-caprolactone), and ethylene vinyl acetate of molecular weights between about 10,000 and 80,000. About 0 to about 100 weight percent polycaprolactone and about 100 to about 0 weight percent of the ethylene vinyl acetate are used based on the total weight of the polymeric material and, preferably, about 50 % each of polycaprolactone and ethylene vinyl acetate is used. The polymeric material used is preferably greater than about 99 % pure and the active agents is preferably greater than about 97 % pure. One of ordinary skill in the art will recognize that in compounding, the conditions under which compounding is carried out will need to take into account the characteristics of the active agent to ensure that the active agents do not become degraded by the process. The polycaprolactone and ethylene vinyl acetate preferably are combined with the desired active agent or agents, micro-compounded, and then extruded as a fiber. The fibers are then cut to the desired length and inserted into the reservoir through one or more plug openings.

20

The amount of active agent used in the plugs of the invention will depend upon the active agent or agents selected, the desired doses to be delivered via the punctual plug, the desired release rate, and the melting points of the active agent and active agent-containing material. Preferably, the amount used is a therapeutically effective amount meaning an amount effective to achieve the desired treatment, inhibitory, or prevention effect. Typically, amounts of about 0.05 to about 8,000 micrograms of active agents may be used.

25

In certain aspects of the invention, the reservoir can be refilled with a material after substantially all of the active agent-containing material has dissolved or degraded and the active agent is released. For example, the new active agent-

30

containing material can be the same as, or different from, the previous polymeric material, and can contain at least one active agent that is the same as, or different from the previous active agent. Certain punctal plugs used for particular applications can preferably be refilled with a material while the punctal plugs remain inserted in the lacrimal canaliculus, while other punctal plugs are typically removed from the lacrimal canaliculus, a new material is added, and the punctal plugs are then reinserted into the lacrimal canaliculus.

When the active agent-containing material is combined with the active agent, the material may also contain one or more materials that are insoluble in water and non-biodegradable, but from which the active agent can diffuse. For example, if the material is a polymeric material, the material may be composed of one or more polymers that are insoluble in water and non-biodegradable. Suitable polymers of this type include, for example, cross-linked polymers, such as, for example, cross-linked poly(ethylene glycol), poly(ethylene oxide), poly(propylene glycol), poly(vinyl alcohol), poly(hydroxyethyl methacrylate), poly(vinylpyrrolidone), polyacrylic acid, poly(ethyloxazoline), and poly(dimethyl acrylamide). These polymers can be copolymerized or blended with one or both of hydrophobic polymers and monomers. Additional polymers that are insoluble in water and non-biodegradable include, without limitation, silicone; silicone blends; silicone copolymers including, without limitation, hydrophilic monomers of pHEMA, polyethylene glycol, polyvinylpyrrolidone, and glycerol; silicone hydrogel polymers such as, for example, those described in U.S. Patent Nos. 5,962,548, 6,020,445, 6,099,852, 6,367,929, and 6,822,016, incorporated herein in their entireties by reference; phospholipids including, without limitation, phosphoryl choline derivatives; polysulfobetains; polysaccharides and carbohydrates including, without limitation, hyaluronic acid, dextran, hydroxyethyl cellulose, hydroxyl propyl cellulose, gellan gum, guar gum, heparan sulfate, chondroitin sulfate, and heparin; proteins including, without limitation, albumin and ovalbumin; polyamino acids; fluorinated polymers including, without limitation, PTFE, PVDF, and teflon; polypropylene;

polyethylene; nylon; and EVA. Additional examples of suitable polymers that are either or both insoluble in water and non-biodegradable include, without limitation,
5 silicones, polyurethanes, cyanoacrylates, and polyacrylic acid.

The punctal plugs described herein can be used to deliver various active agents for the one or more of the treatment, inhibition, and prevention of numerous diseases and disorders. Each punctal plug can be used to deliver at least one active
10 agent and can be used to deliver different types of active agents. For example, the punctal plugs can be used to deliver azelastine HCl, emadastine difumerate, epinastine HCl, ketotifen fumarate, levocabastine HCl, olopatadine HCl, pheniramine maleate, and antazoline phosphate for one or more of the treatment, inhibition, and prevention of allergies. The punctal plugs can be used to deliver
15 mast cell stabilizers, such as, for example, cromolyn sodium, lodoxamide tromethamine, nedocromil sodium, and permirolast potassium.

After the plugs is filled with the active agent, the plug is sterilized by any convenient method including, without limitation, ethylene oxide, autoclaving,
20 irradiation, and the like and combination thereof. Preferably, sterilization is carried out through gamma radiation or use of ethylene oxide.

The punctal plugs can be used to deliver mydriatics and cycloplegics including, without limitation, atropine sulfate, homatropine, scopolamine HBr,
25 cyclopentolate HCl, tropicamide, and phenylephrine HCl. The punctal plugs can be used to deliver ophthalmic dyes including, without limitation, rose bengal, sissamine green, indocyanine green, fluorexon, and fluorescein.

The punctal plugs can be used to deliver corticosteroids including, without
30 limitation, dexamethasone sodium phosphate, dexamethasone, fluoromethalone,

fluoromethalone acetate, loteprednol etabonate, prednisolone acetate, prednisolone sodium phosphate, medrysone, rimexolone, and fluocinolone acetonide. The punctal plugs can be used to deliver non-steroidal anti-inflammatory agents including, without limitation, flurbiprofen sodium, suprofen, diclofenac sodium, ketorolac tromethamine, cyclosporine, rapamycin methotrexate, azathioprine, and bromocriptine.

The punctal plugs can be used to deliver anti-infective agents including, without limitation, tobramycin, moxifloxacin, ofloxacin, gatifloxacin, ciprofloxacin, gentamicin, sulfisoxazole diolamine, sodium sulfacetamide, vancomycin, polymyxin B, amikacin, norfloxacin, levofloxacin, sulfisoxazole diolamine, sodium sulfacetamide tetracycline, doxycycline, dicloxacillin, cephalixin, amoxicillin/clavulante, ceftriaxone, cefixime, erythromycin, ofloxacin, azithromycin, gentamycin, sulfadiazine, and pyrimethamine.

The punctal plugs can be used to deliver agents for the one or more of the treatment, inhibition, and prevention of glaucoma including, without limitation, epinephrines, including, for example: dipivefrin; alpha-2 adrenergic receptors, including, for example, aproclonidine and brimonidine; betablockers including, without limitation, betaxolol, carteolol, levobunolol, metipranolol, and timolol; direct miotics, including, for example, carbachol and pilocarpine; cholinesterase inhibitors, including, without limitation, physostigmine and echothiophate; carbonic anhydrase inhibitors, including, for example, acetazolamide, brinzolamide, dorzolamide, and methazolamide; prostoglandins and prostamides including, without limitation, latanoprost, bimatoprost, uravoprost, and unoprostone cidofovir.

The punctal plugs can be used to deliver antiviral agents, including, without limitation, fomivirsen sodium, foscarnet sodium, ganciclovir sodium, valganciclovir

HCl, trifluridine, acyclovir, and famciclovir. The punctal plugs can be used to deliver local anesthetics, including, without limitation, tetracaine HCl, proparacaine HCl, proparacaine HCl and fluorescein sodium, benoxinate and fluorescein sodium, and benoxinate and fluorexon disodium. The punctal plugs can be used to deliver antifungal agents, including, for example, fluconazole, flucytosine, amphotericin B, itraconazole, and ketocaonazole.

The punctal plugs can be used to deliver analgesics including, without limitation, acetaminophen and codeine, acetaminophen and hydrocodone, acetaminophen, ketorolac, ibuprofen, and tramadol. The punctal plugs can be used to deliver vasoconstrictors including, without limitation, ephedrine hydrochloride, naphazoline hydrochloride, phenylephrine hydrochloride, tetrahydrozoline hydrochloride, and oxymetazoline. Finally, the punctal plugs can be used to deliver vitamins, antioxidants, and nutraceuticals including, without limitation, vitamins A, D, and E, lutein, taurine, glutathione, zeaxanthin, fatty acids and the like.

The active agents delivered by the punctal plugs can be formulated to contain excipients including, without limitation, synthetic and natural polymers, including, for example, polyvinylalcohol, polyethyleneglycol, PAA (polyacrylic acid), hydroxymethyl cellulose, glycerine, hypromelos, polyvinylpyrrolidone, carbopol, propyleneglycol, hydroxypropyl guar, glucam-20, hydroxypropyl cellulose, sorbitol, dextrose, polysorbate, mannitol, dextran, modified polysaccharides and gums, phospholipids, and sulphobetains.

The invention will be clarified further by consideration of the following, non-limiting examples.

Examples

Example 1

- 5 A 1.50 g amount of epsilon polycaprolactone with an average M_w of approximately 14,000 and an average M_n of approximately 10,000 by GPC (available from Aldrich) was combined with 1.50 g EVA (EVATANETM, Arkema), and 1.00 g of bimatoprost (Cayman Chemicals), each with a purity of greater than
- 10 approximately 97 %. The mixture was then placed in a twin-screw micro-compounder Model No. 20000 from DACA Industries, Inc. that was fitted with a 0.25 mm die and compounded for 15 min. at 120 rpm and 67° C. Following compounding, the mixture was extruded into fibers at 75° C.
- 15 The fibers were cut into approximately 1.5 mm in length sections and inserted into the opening of Sharpoint ULTRATM plug, available from Surgical Specialties. 70 plugs with drug and 30 placebo each of a 0.6 mm length were made and 70 drugs with drug and 50 placebo each of a 0.8 mm length were made. To
- 20 insert the fiber, each plug was positioned under a stereomicroscope and tweezers were used to insert a fiber into the opening of each of the plugs. Each plug was then placed in a glass vial and packed in a pouch. Gamma radiation sterilization was then performed with a total does of 15 Kgy at about 0.2 Kgy/min.

Example 2

- 25 A plug made according to the method of Example 1 that is 0.6 mm in length and a fiber having a diameter of approximately 0.2 mm and containing 25 % w/w was placed in a glass vial with 1 cc of phosphate buffered saline having a pH of 7.4. The vial was placed in an incubator at 37° C and gently agitated. Aliquots of 1cc were collected at intervals at 3, 8, and 24 hours and then weekly and analyzed for
- 30 drug content via HPLC. Figure 11 is a graph depicting the results of the analysis.

Example 3

A plug made according to the method of Example 1 that is 0.9 mm in length
5 and a fiber with an approximately 0.6 mm diameter and containing 25 % was placed
in a glass vial with 1 cc of phosphate buffered saline having a pH of 7.4. The vial
was placed in an incubator at 37° C and gently agitated. Aliquots of 1 cc were
collected at intervals as set forth in Example 2 and analyzed for drug content via
HPLC. Figure 12 is a graph depicting the results of the analysis.

What is claimed is:

1. A punctal plug, comprising:
a body having a first end, a second end, and a lateral surface extending
5 between the two ends;
a reservoir contained within the body, wherein the reservoir comprises at least one opening, and contains an active agent-containing material that comprises at least one active agent; and
wherein the body is impermeable to the active agent.
- 10 2. A punctal plug, comprising:
a body having a first end, a second end, and a lateral surface extending between the two ends;
a reservoir contained within the body, wherein the reservoir comprises an opening at the first end, the second end, or both the first and second ends of the
15 body, and contains an active agent-containing material that comprises at least one active agent; and
wherein the body is impermeable to the active agent.
3. The punctal plug of claim 1, wherein the lateral surface of the body has an outer diameter that is substantially circular in shape and a portion of the lateral
20 surface has an outer diameter that is greater than that of the remainder of the lateral surface
4. The punctal plug of claim 2, wherein the lateral surface of the body has an outer diameter that is substantially circular in shape and a portion of the lateral
surface has an outer diameter that is greater than that of the remainder of the lateral
25 surface

5. The punctal plug of claim 1, 2, 3, or 4, wherein the active agent-containing material is at least partially water-soluble, dissolves over time, and releases the active agent through an opening in the reservoir as it dissolves.
6. The punctal plug of claim 1, 2, 3, or 4, wherein the active agent-containing material is biodegradable, degrades over time, and releases the active agent through an opening in the reservoir as it degrades.
7. The punctal plug of claim 1, 2, 3, or 4, wherein the active agent-containing material is insoluble in water and non-biodegradable and the active agent passively diffuses from the material through an opening in the reservoir.
8. The punctal plug of claim 3, or 4, wherein the portion of the lateral surface that has an outer diameter greater than that of the remainder of the lateral surface secures the punctal plug in a lacrimal canaliculus when the punctal plug is inserted in the lacrimal canaliculus.
9. The punctal plug of claim 1, 2, 3, or 4, wherein when the punctal plug is inserted into a lacrimal canaliculus of an eye, an opening in the reservoir faces the eye and the active agent is released into a tear fluid of the eye.
10. The punctal plug of claim 1, 2, 3, or 4, wherein when the punctal plug is inserted into a lacrimal canaliculus of an eye, an opening in the reservoir faces a nasolacrimal duct and the active agent is released into the nasolacrimal duct.
11. The punctal plug of claim 1, 2, 3, or 4, wherein when the punctal plug is inserted into a lacrimal canaliculus of an eye, an opening in the reservoir faces the eye and the active agent is released into a tear fluid of the eye and another opening in the reservoir faces a nasolacrimal duct and the active agent is also released into the nasolacrimal duct.

12. A punctal plug for the delivery of active agent to the tear fluid of an eye and to a nasolacrimal duct, comprising:

5 a body comprised of a flexible polymeric material that conforms to the shape of a lacrimal canaliculus when the punctal plug is inserted into the lacrimal canaliculus;

a reservoir contained within the body, wherein the reservoir comprises at least one opening and contains an active agent-containing material that comprises at least one active agent, and the body is impermeable to the active agent; and

10 a collarette that rests on the exterior of the lacrimal punctum when the punctal plug is inserted into the lacrimal canaliculus.

13. The punctal plug of claim 12, wherein the active agent-containing material is at least partially water-soluble, dissolves over time, and releases the active agent through an opening in the reservoir as it dissolves.

15 14. The punctal plug of claim 13, wherein the reservoir is refilled with a active agent-containing material comprising at least one active agent that is the same as or different from the previous active agent after substantially all of the previous active agent-containing material has dissolved.

20 15. The punctal plug of claim 19, wherein the active agent-containing material is biodegradable, degrades over time, and releases the active agent through an opening in the reservoir as it degrades.

16. The punctal plug of claim 13, wherein the reservoir is refilled with an active agent-containing material comprising at least one active agent that is the same as or

different from the previous active agent after substantially all of the previous material has degraded.

17. The punctal plug of claim 12, wherein the active agent-containing material is
5 insoluble in water and non-biodegradable and the active agent passively diffuses from the material through an opening in the reservoir.

18. The punctal plug of claim 12, wherein the reservoir and the body are
coterminous.

10

19. The punctal plug of claim 12, wherein when the punctal plug is inserted into the lacrimal canaliculus an opening in the reservoir faces the eye and the active agent is released into the tear fluid of the eye.

15 20. The punctal plug of claim 12, wherein when the punctal plug is inserted into the lacrimal canaliculus an opening in the reservoir faces the nasolacrimal duct and the active agent is released into the nasolacrimal duct.

21. The punctal plug of claim 12, wherein when the punctal plug is inserted into
20 the lacrimal canaliculus an opening in the reservoir faces the eye and the active agent is released into the tear fluid of the eye and another opening in the reservoir faces the nasolacrimal duct and the active agent is also released into the nasolacrimal duct.

22. A punctal plug, comprising:
25 a body having a first end, a second end, and a lateral surface extending between the two ends;

a reservoir contained within the body, wherein the reservoir comprises at least one opening, and contains an active agent-containing material that is a fiber or
5 fiber-like structure comprising at least one active agent; and
wherein the body is impermeable to the active agent.

23. The punctual plug of claim 22, wherein the lateral surface of the body has an outer diameter that is substantially circular in shape and a portion of the lateral surface has an outer diameter that is greater than that of the remainder of the lateral
10 surface

24. The punctual plug of claim 22, wherein the reservoir contains a fiber that comprises about 0 to about 100 weight percent of polycaprolactone and about 100 to about 0 weight percent of ethylene vinyl acetate.

25. The punctual plug of claim 22, wherein the reservoir contains a fiber that
15 comprises about 50 weight percent of polycaprolactone and about 50 weight percent of ethylene vinyl acetate.

1/6

FIG. 1




FIG. 2





FIG. 3



2/6

FIG. 4




FIG. 5





FIG. 6



3/6

FIG. 7




FIG. 8






FIG. 9




4/6

FIG. 10



5/6

FIG. 11

6/6

FIG. 12